

SPECTROSCOPIC AND THEORETICAL CONSIDERATIONS
OF CYCLOPROPENONE

BY

Caroline P. Starrs, G.R.S.C., Ireland.

A Thesis
submitted to the Department of Chemistry
in partial fulfillment of the requirements
for the degree of
Master of Science

August 1988
Brock University
St. Catharines, Ontario.

graduated Fall 1988

(C) Caroline P. Starrs, 1988

ABSTRACT

The absorption spectrum of cyclopropenone has been photographically recorded under conditions of long pathlength (60 m), and moderate resolution (7.8 \AA/mm). The absorption was assigned to the electron promotion, $n \rightarrow \pi^*$ (C=O), and to the electronic transition $\tilde{\text{B}}^1\text{A}_2 \rightarrow \tilde{\text{X}}^1\text{A}_1$. The spectrum proved to be limited in band structure, which hindered the vibrational assignment. Structures and vibrational frequencies were calculated for the $\tilde{\text{X}}^1\text{A}_1$, $\tilde{\text{a}}^3\text{B}_1$, and $\tilde{\text{b}}^3\text{A}_2$ states at the 6-31g* Hartree-Fock level. The calculations confirmed the assigned transition and predicted the position and structure of the second $\tilde{\text{a}}^3\text{B}_1$ system.